

Higher Harmonics in Non-Linear Vacuum from QED Effects Without Low Mass Intermediate Particles

J. Tito Mendonça¹⁾, J. Dias de Deus²⁾, and P. Castelo Ferreira²⁾

¹⁾ CFP and CFIF, Instituto Superior Técnico, 1096 Lisboa, Portugal and

²⁾ CENTRA, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

(Dated: February 2, 2008)

We show that in the presence of a slowly rotating strong transverse magnetic field there is an infinite spectrum of harmonic wave functions A_n due to the first order QED correction (in α^2) given by the Euler-Heisenberg Lagrangian. The frequency shifts are integer multiples $\pm\omega_0 n$ of the magnetic field angular frequency rotation $\omega_0 = 2\pi\nu_m$ and the several modes n are coupled to the nearest harmonics $n \pm 1$. This is a new effect due to QED vacuum fluctuations, not exploit before, that can explain, both qualitatively and quantitatively, the recent experimental results of the PVLAS collaboration without the need of a low mass intermediate particle, hence may dismiss the recent claim of the discovery of the axion.

In quantum field theory the vacuum is not empty, in the sense that virtual particle pair creation and annihilation are taking place. Considering the case of QED -Quantum Electrodynamics- in the presence of external electromagnetic fields the vacuum shows properties close to the properties of any other optical medium, as dichroism and birefringence. These effects can be considered as first order corrections to the Maxwell theory [1, 2, 3, 4], and the basic diagrams, correcting the photon propagator, are shown in figure 1. In the presence of low mass neutral particles coupling to two photons, like the π^0 and, more interesting, the axion -Primakov effect- or the graviton, there are additional non-linear effects (see figure 1), which may be useful in directly detect the axion [5, 6].

In this letter we present and exploit an effect not considered before, in vacuum a strong rotating magnetic field generates an infinite set of higher harmonics. We show that recent experimental results [7] can be explained in the basis of pure QED which dismisses the recent claim of the discovery of the axion [10].

In the QED lowest order correction (radiative corrections in α^2) to the Maxwell Lagrangian $\mathcal{L}_{\text{Maxwell}} = -F_{\mu\nu}F^{\mu\nu}$ is given by the regularized Euler-Heisenberg Lagrangian $\mathcal{L}^{(2)}$ [1, 2]

$$\mathcal{L}^{(2)} = \xi \left[4 (F_{\mu\nu}F^{\mu\nu})^2 + 7 (\epsilon^{\mu\nu\delta\rho} F_{\mu\nu}F_{\delta\rho})^2 \right] \quad (1)$$

where $\xi = 2\alpha/(45B_c^2) \sim \alpha^2$ and the critical field is $B_c = m_e^2 c^2/(e\hbar) = 4.414 \times 10^9 \text{ T}$ [4]. The usual procedure is to take a decomposition of the gauge connection F into an external component F_0 plus an internal component $f_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, such that we obtain $F_{\mu\nu} = F_{0,\mu\nu} + f_{\mu\nu}$, and consider the lowest order diagrams of photon interactions due to the background electromagnetic fields. There are two main processes that may be considered, *photon splitting* and *photon dispersion*. Depending on the practical application and experimental setups there are other processes, such as, *photon-photon interactions* [8] and other photon alternative processes [9] that are relevant. In this work we address only the process of *photon dispersion* in the low frequency and weak field limit, i.e. $\omega \ll 2m_e c^2/\hbar$ and $|F| \ll m_e c^2/(\hbar e)$.




FIG. 1: Feynman diagrams corresponding to the electron loop that contributes to the photon dispersion effect and the axion exchange.

The relevant diagrams are the ones containing one internal fermion loop with two external photons and an even number of exchanged photons between the fermion loop and the external electromagnetic field (see figure 1).

We obtain that the relevant radiative corrections to the usual classical wave equation in order α^2 is linear in the photon field A [4]

$$(\nabla^2 - \frac{1}{c^2} \partial_t^2) \mathbf{A} = \Lambda \cdot \mathbf{A} + O(\alpha^4), \quad (2)$$

where the matrix Λ has the eigenvalues [3, 4]

$$\lambda_{\parallel} = 14\xi(\omega/c)^2 |\mathbf{Q}|^2 \text{ and } \lambda_{\perp} = 8\xi(\omega/c)^2 |\mathbf{Q}|^2, \quad (3)$$

where $\mathbf{Q} = \mathbf{k} \times \mathbf{E}_0/|\mathbf{k}| + \mathbf{k} \times (\mathbf{k} \times \mathbf{B}_0)/|\mathbf{k}|^2$ and $\mathbf{k}$ stands for the light wave vector. $\mathbf{E}_0$ and $\mathbf{B}_0$ are the external fields and the directions $\parallel$ and $\perp$ correspond respectively

to the parallel and transverse directions to the external magnetic field.

We consider linear polarized light traveling in vacuum in the z direction with wave number $\mathbf{k}$ and frequency ω under a slowly rotating transverse magnetic field rotating with angular frequency $\omega_0 = 2\pi\nu_m \ll \omega$ such that we have $\mathbf{A} = (A_x, A_y, 0)$, $\mathbf{k} = (0, 0, k)$, $w = ck$, $\mathbf{B}_0(t) = B_0 [\sin(\omega_0 t), \cos(\omega_0 t), 0]$, $\mathbf{E}_0 = 0$ and $|\mathbf{Q}| = cB_0$. In these conditions the gauge field depends only on z and we can write the equation (2) for $\mathbf{A}$ in terms of the parallel direction ($\parallel$) and transverse direction ($\perp$) as derived by Adler [4]. However these directions are rotating with an angular frequency ω_0 , therefore we take the directions of

the decomposition of the fields to be the directions of the magnetic field at some initial time $t = 0$. Then the eigenvalues of the matrix Λ become time dependent since they are valid with respect to the magnetic field direction such that we have

$$\begin{aligned}\Lambda_{\parallel}(t) &= \lambda_{\parallel} \cos(\omega_0 t) - \lambda_{\perp} \sin(\omega_0 t) \\ \Lambda_{\perp}(t) &= \lambda_{\parallel} \sin(\omega_0 t) + \lambda_{\perp} \cos(\omega_0 t)\end{aligned}\quad (4)$$

with $\lambda_{\parallel}$ and $\lambda_{\perp}$ given in (3). Considering the above eigenvalues we obtain the field equation

$$(\partial_z^2 - \frac{1}{c^2} \partial_t^2)(A_{\parallel}^0, A_{\perp}^0) = (\lambda_{\parallel} \cos(\omega_0 t) - \lambda_{\perp} \sin(\omega_0 t))(A_{\parallel}^0, 0) + (\lambda_{\parallel} \sin(\omega_0 t) + \lambda_{\perp} \cos(\omega_0 t))(0, A_{\perp}^0), \quad (5)$$

where the $A_{\parallel}^0$ and $A_{\perp}^0$ correspond to the decomposition of the field along the directions parallel ($\parallel$) and transverse ($\perp$) to the magnetic field at the initial time $t = 0$. For constant magnetic field ($\omega_0 = 0$) these equations imply that the vacuum acquires a birefringence due to the existence of two different refractive indices $\mathbf{n}_i = 1 - \lambda_i/2$ along both the directions $i = \parallel, \perp$. For slowly rotating magnetic fields the usual procedure is to take the zeroth order expansion of the cosines and sines that hold the same effect [6]. However we show next that even for small ω_0 equation (5) implies, not just a vacuum birefringence, but instead a coupling between the neighbor harmonics separated by frequency shifts of $\pm\omega_0$. It is important to note that equation (5) is valid for slow rotating external fields (see [4] for further details), i.e. $\omega_0 \ll \omega$.

We can cast equation (5) in complex form by using the definitions $\lambda_{\pm} = \lambda_{\parallel} \pm i\lambda_{\perp}$, $A_{\pm} = A_{\parallel} \pm iA_{\perp}$ and the usual relations for the trigonometric functions and consider a mode decomposition of the form

$$\begin{aligned}A_{\pm}(z, t) &= \sum_{n=-\infty}^{+\infty} A_n^{\pm}(t) e^{i[kz - \omega_n t + \theta_n^{\pm}]}, \\ ck_n = \omega_n &= \omega + \omega_0 n, \quad \theta_n^{\pm} = \left(-\theta_{\lambda} \pm \frac{\pi}{4}\right) n, \\ \lambda_0 &= \sqrt{\lambda_{\parallel}^2 + \lambda_{\perp}^2}, \quad \theta_{\lambda} = \arctan\left(\frac{\lambda_{\perp}}{\lambda_{\parallel}}\right).\end{aligned}\quad (6)$$

Given the above mode decomposition the differential equation becomes a recursion relation in the several frequency modes A_n

$$i \frac{k}{\sqrt{2} c \lambda_0} \dot{A}_n = A_{n+1} + A_{n-1} \quad (7)$$

where $A_n = A_n^+ = e^{-i\pi/2} A_n^-$. We neglected a second derivative term $\ddot{A}_n$ and considered a dispersion relation

that renders a tower of refractive indices in each direction. This is the same approach of [4] in consistent field approach. Here we have for each mode n and each direction $i = \parallel, \perp$ a different refractive index

$$kc = \mathbf{n}_i^n(\omega) \omega, \quad \mathbf{n}_i^n(\omega) = 1 + \frac{\omega_0}{\omega} n. \quad (8)$$

The solutions up to an overall phase for the recursive relation (7) are modified Bessel function of the first kind

$$A_n(t) = (-1)^n I_n(i\tau), \quad \tau = \frac{\sqrt{2} c \lambda_0 t}{k}. \quad (9)$$

We note that indeed the second derivative of A_n is of order α^4 and can be safely neglected. These functions are normalized for any τ and for even n the functions are pure reals while for odd n the functions are pure imaginary. The only mode that is non null at $\tau = 0$ is $I_0(i\tau)$ such that only for larger values of t the amplitudes of the other modes increase. These properties are important when imposing the initial conditions, i.e. matching our solution with the incident wave at the initial time $t = 0$.

The results obtain so far are very interesting and describe both vacuum birefringence and a generalized dichroism in the presence of a strong rotating magnetic field. The tower of refractive indices (8) correspond to a birefringence effect quantized in terms of the quantity ω_0/ω . The Bessel functions solutions account for a generalized dichroism effect that decreases the amplitude of the incident wave in one mode and increases it in the nearest modes, we note that the QED effects considered here do preserve momenta such that there is no real loss of energy, it simply is transferred from one mode to the nearest modes. These effects can be understood as follows, from equation (7) we see that the variation of each mode amplitude A_n over time contributes to the nearest modes $A_{n\pm 1}$, this accounts for the generalized dichroism effect in the sense that no real absorption happens,

therefore we have a *true* rotation of polarization (in the sense considered in [7]). As for birefringence we note that by combining the nearest modes (e.g. $n_{\parallel} = n$ and $n_{\perp} = n \pm 1$) in different directions we will have a net birefringence effect given by $\mathbf{n}_{\perp} - \mathbf{n}_{\parallel} = \pm \omega_0/\omega$ such that each mode combination correspond to a wave with elliptic polarization. It is also interesting to note that, although the description for the propagating wave is given in terms

of the collective field A , we may as well interpret these solutions as cross section for photons. In such interpretation the allowed configurations for a single photon are independent in both spatial dimensions ($\parallel, \perp$), such that indeed we have elliptic polarized waves.

A solution for the electric field compatible with linearized polarized waves at the initial time $\tau = 0$ is

$$\mathbf{E} = E_0 \left\{ \sum_{n=-\infty}^{+\infty} [I_{2n}(i\tau) \cos(kz - \omega_{2n}t + \theta_{2n}) - i I_{2n-1}(i\tau) \sin(kz - \omega_{2n-1}t - \theta_{2n-1})] \right\} (\sin(\theta_0), \cos(\theta_0)) \quad (10)$$

Here E_0 is the electric field amplitude, $\omega_n = \omega + \omega_0 n$ and $\theta_n = -(\theta_{\lambda} + \pi/4)n$. This field is defined in the referential of the parallel and transverse direction to the magnetic field at the initial time $t = 0$ such that θ_0 stands for the angle between the original linear polarization direction and the magnetic field direction at $t = 0$.

Take the $n_{\parallel} = n_1$ mode in one of the directions together with one of the other modes $n_{\perp} = n_2 > n_1$ in the other direction such that this combination has now an elliptic polarization and the difference of argument of the trigonometric functions corresponding to each direction correspond to a phase difference of $\Delta\varphi_{n_2-n_1} = (n_2 - n_1)(\omega_0 t + \theta_{\lambda} + \pi/4)$.

We proceed to compare our results with the recent experimental results of the PVLAS collaboration [7]. For the apparatus of that experiment an incident polarized laser beam traverses a region under a strong rotating magnetic field such that it is possible to measure polarization rotations (that correspond to vacuum dichroism) and ellipticities (that correspond to vacuum birefringence).

We can write the amplitude measured in the experience for our solutions as

$$\mathcal{I} = \left[2 \sum_{n=1}^{+\infty} \Gamma_n |I_n(i\tau)| \cos(n(\omega_0 t + \theta_{\lambda} + \theta_0 + \pi/4)) + \eta_0 \cos(\omega_S t + \theta_S) + \Gamma_0 \right]^2 \quad (11)$$

Here $\eta_0 \cos(\omega_S t + \theta_S)$ stands for the modulator wave [7] and heterodyne detection is used to identify the several harmonics present due to rotations. Γ_n are corrections that encode uncompensated dispersions for the several modes n due, either to systematic deviations, or unaccounted processes. In (11) we sum over rotations in one direction only, that is why we have a factor of 2 in front of the sum. By expanding the square of this expression we obtain the angular frequency spectrum and the respective amplitudes and phases for each of them. We list the more significant frequencies that can be measured near

Frequency	$\mathcal{I}$	Phase
0	$\Gamma_0^2 + 2 \sum_{n=1}^{+\infty} (\Gamma_n I_n)^2 + \frac{\eta_0^2}{2}$	--
ω_S	$2\Gamma_0 \eta_0$	θ_S
$\omega_S \pm \omega_0 n$	$2\Gamma_n I_n \eta_0$	$\theta_S \pm n(\theta_{\lambda} + \theta_0 + \pi/4)$

TABLE I: The relevant angular frequency spectrum measured near the modulator frequency ω_S with respective amplitudes and phases. There is an infinite set of other frequencies that have lower amplitudes.

the modulator frequency ω_S and list them in table I.

We take the values of figure 2 of the original reference [7] (expressed in I.S. units)

$$\omega = 1.772 \times 10^{+15} \text{ s}^{-1} \quad \omega_0 = 0.7\pi \text{ s}^{-1} \quad (12)$$

$$\Delta z = c\Delta t = 4.6 \times 10^{+4} \text{ m} \quad B_0 = 5.5 \text{ T}.$$

The theoretical results obtained above are valid for these experimental conditions, in particular the magnetic field is slowly rotating ($\omega_0 \ll \omega$) and obey the low frequency and weak field approximations, respectively $\omega \sim 10^{15} \ll 2m_e c^2/\hbar \sim 10^{21}$ and $B_0^2 \sim 10 \ll m_e c^2/(e\hbar) \sim 10^{39}$. For the values given in (12) and using the finesse of the Fabry-Perot cavity $\mathcal{F} \approx 7 \times 10^4$ (corresponding to an increase of optical path $N \approx 4.46 \times 10^4$) [7] we have from (9) with $\tau = 0.0088$ [3] that $|I_0(i\tau)| \approx 0.9998$, $|I_1(i\tau)| \approx 4.384 \times 10^{-3}$, $|I_2(i\tau)| \approx 9.609 \times 10^{-6}$ and $|I_3(i\tau)| \approx 1.404 \times 10^{-8}$. We use these values to fit our solutions to the data expressed in figure 2 of reference [7]. We considered that the background noise is $\approx -135 \text{ dBVrms}$ and assumed values for η_0 and the Γ_i 's that allow us to fit the experimental results. These values are given as an example and are not based in a detailed analysis. In table II we list a possible set of values (in dBVrms) that fits the experimental results of [7]. We note that the value obtained for the mode $n = 3$ ($\approx -145.65 \text{ dBVrms}$) is

frequency (Hz)	$\mathcal{I}_{\text{exp}}$ (dBVrms)	$\eta_0 I_n$ (dBVrms)	Γ_i
506	≈ -51	--	$\approx +16.93$
506 ± 0.35	≈ -91	≈ -90.70	$\approx +0.704$
506 ± 0.70	≈ -118	≈ -117.30	≈ -0.704
506 ± 1.05	< -135	≈ -145.65	--

TABLE II: We list the experimental values taken from figure 2 of reference [7] with the theoretical values given by the $I_n(i\tau)$ up to $n = 3$. We considered $\eta_0 \approx -68.63$ dBVrms. This value, as well as the value of the Γ_n 's, are only given as an example that it is possible to fit the data to our solutions. The theoretical value found for the amplitude of a rotation corresponding to $n = 3$ is ≈ -145.65 dBVrms and is therefore below the noise level of the experiment that we considered (≈ -135 dBVrms).

below the noise level (≈ -135 dBVrms), hence it could not be detected for the available data.

To conclude we compute the expected rotation values corresponding to the second harmonic $n = 2$. By direct geometrical analysis we conclude that the angular rotations of polarization are given by the generic expression $\varphi = \theta_0 - \arctan\left(\frac{E_{\perp}^{\text{out}}}{E_{\parallel}^{\text{out}}}\right)$, where $\mathbf{E}^{\text{out}} = (E_{\parallel}^{\text{out}}, E_{\perp}^{\text{out}})$ stands for the radiation electric field reaching the detector. For a given harmonic n we have the possible combinations $\mathbf{E}_n^{\text{out}+} = [I_0 \sin(\theta_0) \cos(\delta(t, z)), I_n \cos(\theta_0) \cos(\delta(t, z) + \omega_0 \Delta t + \arctan(7/4) + \pi/4)]$ and $\mathbf{E}_n^{\text{out}-} = [I_n \cos(\theta_0) \cos(\delta(t, z) + \omega_0 \Delta t + \arctan(7/4) + \pi/4), I_0 \sin(\theta_0) \cos(\delta(t, z))]$. Here the phase $\delta(t, z) = kz - \omega t$ is the phase of the wave when reaching the detector and the sine and cosine of θ_0 is due to the angle between the original polarization and the magnetic field when the radiation enters the magnetic field as expressed in the solution (10).

Based in the values (12) with $\theta_0 = \pi/4$ (that corre-

sponds to the maximum effect) and assuming $\delta(t, z) = 0$ we obtain for the rotations corresponding to $n=2$ $\varphi_{n=2}^+ \approx 3.78 \times 10^{-6}$ rad. This value is one order of magnitude higher than the experimental values measured ($\sim 10^{-7}$). A possible explanation for this result may be due to a different initial value of the angle between the polarization and the magnetic field, θ_0 . If we consider for instance $\theta_0 = \pi/48$ we obtain $\varphi_{n=2}^+ \approx 2.48 \times 10^{-7}$ rad which is closer to the experimental observed rotation. Other possibilities may be due to an effective lower path length due either to decoherence or unaccounted losses.

So we have presented a new effect of vacuum polarization rotation due to external strong magnetic fields based in pure QED. This effect generates an infinite number of harmonics and explains, both qualitatively and quantitatively, the PVLAS experimental results [7]. These experimental results show that the PVLAS experiment gives (as far as we know for the first time) a beautiful demonstration of the QED properties of vacuum and have before hand registered an effect that have not been justified at theoretical level. Our results may dismiss the recent claim of the finding of the axion [10]. Both for future use in experiments currently being planned and for a correct interpretation of already detected (and to be detected) physical phenomena [11], a full analysis of existent data is required. In particular, we note that, if the noise level can be decreased, the third peak ($n = \pm 3$) should be detected.

Erratum

A new mechanism was proposed to explain the observed spectrum in the PVLAS experiment [7]. We have predicted a series of new signals with frequencies $\omega + n\omega_0$, for n integer, where ω is the laser frequency and ω_0 the angular frequency of the rotating magnetic field. But two mistakes were made in this work. First, the secondary signals have frequencies $\omega + 2n\omega_0$. Second, our numerical estimates are wrong by a factor of 10^3 [12], and therefore cannot explain the observed signal. It should be stressed that signals at frequencies $\omega + n\omega_0$ can still be generated, but they are not relevant to the PVLAS experiment.

Acknowledgements

Work of PCF supported by SFRH/BPD/17683/2004.

-
- | | |
|---|---|
| <p>[1] W. Heisenberg and H. Euler, Z. Physik 98 (1936) 714, physics/0605038.</p> <p>[2] J. Schwinger, Phys. Rev. 82 (1951) 664.</p> <p>[3] Z. Bialynicka-Birula and I. Bialynicka-Birula, Phys. Rev. D2 (1970) 2341.</p> <p>[4] S. L. Adler, Ann. Phys. 67 (1971) 559.</p> <p>[5] L. Maiani, R. Petronzio and E. Zavattini, Phys. Lett. B175 (1986) 359.</p> <p>[6] G. Raffelt and L. Stodolski, Phys. Rev. D37 (1988) 1237.</p> <p>[7] E. Zavattini and al., Phys. Rev. Lett. 96 (2006) 110406.</p> <p>[8] E. Lundström et al., Phys. Rev. Lett. 96 (2006) 083602.</p> | <p>[9] M. Marklund, P. K. Shukla, Rev. Mod. Phys. 78 (2006) 591.</p> <p>[10] S. Lamoreaux, Nature Vol. 441 (2006) 31.</p> <p>[11] E. Massó, hep-ph/0607215; Y. N. Gnedin, M.Y. Piotrovich, and T. M. Natsvlshvili, astro-ph/0607294; U. Kotz, A. Ringwald, and T. Tschentscher, hep-ex/0606058; E. Massó and J. Redondo, hep-ph/0606163; A.V. Afanasev, O. K. Baker, and K.W. McFarlane, hep-ph/0605250.</p> <p>[12] S. L. Adler, hep-ph/0611267.</p> |
|---|---|